

AMENDMENTS TO THE CLAIMS

This listing of claims will replace all prior versions, and listings, of claims in the application:

LISTING OF CLAIMS:

1. (Currently Amended) A method of enumerating the number of CD4+ lymphocytes in a cell sample, comprising:
 - a) determining the absolute number of CD45+ cells in a cell sample ~~by means of using a hematology analyzer having a facility for fluorescence measurement or via a flow cytometer;~~
 - b) determining the proportion of CD45+ cells which are CD4+ lymphocytes using fluorescence measurement via flow cytometry by:
 - (i) establishing a primary ~~gate, defining all discriminator to discriminate~~ CD45+ cells of interest;
 - (ii) establishing a secondary ~~gated discriminator, within said primary gate, defining all discriminator to discriminate~~ CD4+ lymphocytes having CD4^{Bright}/low ~~CD4 Bright low~~ side scatter expression; and
 - (iii) calculating the proportion of CD45+ cells which are CD4+ lymphocytes; and
 - c) multiplying the absolute number of CD45+ cells obtained in step (a), by the proportion of CD45+ cells which are CD4+ lymphocytes, obtained in step (b), thereby determining the absolute number of CD4+ lymphocytes in the cell sample .
2. (Currently Amended) ~~A~~ The method according to claim 1, wherein ~~the number of CD4+ lymphocytes is enumerated in a single platform approach whereby the number of white blood cells counted in step (a) determining the absolute number of CD4+ cells in the cell sample is obtained on a performed by flow cytometer cytometry using a bead-based or volumetric-based counting method and the proportion or percentage of CD4+ lymphocytes determined in step (b) is also performed on a flow cytometer, or whereby the number of white blood cells counted in step (a) is obtained on a hematology analyzer and the proportion~~

~~or percentage of CD4+ lymphocytes determined in step (b) is also performed on a hematology analyzer.~~

3. (Withdrawn) A method according to claim 1, wherein a white blood cell differential is further identified from the total CD45 expressing population identified in step (a) of claim 1, and further comprising the steps of:

- e) (i) determining the percentage of CD4 monocytes as a function of the total CD45 expressing population identified in step (a); and
(ii) calculating the number of CD4 monocytes in the sample by multiplying the percentage of CD4 monocytes determined in step (e(i)) by the number of CD45 cells obtained in step (c);
- f) (i) determining the percentage of CD4 eosinophils as a function of the total CD45 expressing population identified in step (a); and
(ii) calculating the number of CD4 eosinophils in the sample by multiplying the percentage of CD4 eosinophils determined in step (f(i)) by the number of CD45 cells obtained in step (c);
- g) (i) determining the percentage of CD4 negative granulocytes as a function of the total CD45 expressing population identified in step (a); and
(ii) calculating the number of CD4 negative granulocytes in the sample by multiplying the percentage of CD4 negative granulocytes determined in step (g(i)) by the number of CD45 cells obtained in step (c);
- h) (i) determining the percentage of CD4 negative lymphocytes as a function of the total CD45 expressing population identified in step (a); and
(ii) calculating the number of CD4 negative lymphocytes in the sample by multiplying the percentage of CD4 negative lymphocytes determined in step (h(i)) by the number of CD45 cells obtained in step (c);

- i) (i) adding the percentage of CD4 lymphocytes identified in step (b) and the percentage of CD4 negative lymphocytes obtained in step (h(i)) to obtain the percentage of total lymphocytes;
- (ii) determining the percentage of total lymphocytes as a function of the total CD45 expressing population identified in step (a); and
- (iii) calculating the number of total lymphocytes in the sample by multiplying the percentage of total lymphocytes obtained in step (i(ii)) by the number of CD45 cells obtained in step (a).

4. (Withdrawn) A method according to claim 1, further comprising the steps of:

- e) determining the percentage of basophils as a function of the total CD45 expressing population identified in step (a); and
- f) calculating the number of basophils in the sample by multiplying the percentage of basophils determined in step (j) by the number of CD45 cells obtained in step (c).

5. (Previously presented) A method according to claim 1, wherein the sample is whole unlysed blood, unfractionated, fractionated or lysed whole blood.

6. (Withdrawn) A kit including CD4 and CD45 antibodies for use in enumerating the number of CD4 cells in a sample.

7. (Withdrawn) A kit according to claim 6, which further includes instructions for performing the method of enumerating the number of CD 4 cells in a cell sample.

8. (Withdrawn) A kit according to claim 6, which further includes one or more reagents selected from the group consisting of a red cell lysating agent, a stabilizer, a fixative, control cells, media and bead reagents.

9. (Withdrawn) A machine readable medium comprising instructions, which when executed by a machine, cause the machine to perform the method steps of claim 1.

10. (Withdrawn) A machine readable medium according to claim 9, which is configured for use in conjunction with a flow cytometer and/or haematology analyser.

11. (Withdrawn) A machine readable medium according to claim 9, which includes instructions for performing analysis methods selected from the group consisting of impedance, light scatter, fluorescence and precision volume counting.

12. (Withdrawn) The method of claim 1, wherein the method includes the step of enumerating the number of CD4+ lymphocytes or CD4+ T-cells in a cell sample from the patient with HIV or other immune deficiency condition or disease; and wherein the method is used to monitor the immune status of the patient.

13. (Withdrawn) A method according to claim 12, wherein the patient's immune status is determined or monitored to determine the patient's response to antiretroviral treatment.

14. (Currently Amended) ~~A-The~~ method according to claim 2, wherein the ~~single platform determination is performed by bead-based counting method comprises~~ adding a known numbers of beads to the cell sample; and counting the beads and cells simultaneously to obtain the absolute CD45+ cell count.

15. (Currently Amended) ~~A-The~~ method according to claim 1, wherein the number of CD4+ lymphocytes is enumerated in a dual platform approach whereby the number of ~~white blood cells counted~~ CD45+ cells determined in step (a) is obtained ~~from using a~~ hematology analyzer and the proportion ~~or percentage~~ of CD4+ lymphocytes determined in step (b) is performed ~~on using a~~ flow cytometer.

16. (Currently Amended) ~~A-The~~ method according to claim 1, wherein the number of ~~white blood~~ CD45+ cells per volume of blood is determined in step (ea) ~~by determination of total CD45+ expressing cells per volume of blood.~~